



L-Selectin Monoclonal Antibody

Catalog No	YP-mAb-17086
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SELL
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human SELL. AA range:91-140
Specificity	L-Selectin Monoclonal Antibody detects endogenous levels of L-Selectin protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	L-selectin / CD62L;L-selectin;CD62 antigen-like family member L;gp9 MEL;gp9-MEL;SELL;LNHR;LYAM1;Leukocyte adhesion molecule 1;LAM-1;Leukocyte surface antigen Leu-8;Leukocyte-endothelial cell adhesion molecule 1;LECAM1;Lymph node homing receptor;TQ1;gp90-MEL;CD62L
Observed Band	42kD
Calculated Molecular Weight	42kD
Cell Pathway	Cell membrane ; Single-pass type I membrane protein .
Tissue Specificity	Expressed in B-cell lines and T-lymphocytes.
Function	function:Cell surface adhesion protein. Mediates the adherence of lymphocytes to endothelial cells of high endothelial venules in peripheral lymph nodes. Promotes initial tethering and rolling of leukocytes in endothelia.,online information:L-selectin,similarity:Belongs to the selectin/LECAM family.,similarity:Contains 1 C-type lectin domain.,similarity:Contains 1 EGF-like domain.,similarity:Contains 2 Sushi (CCP/SCR) domains.,subunit:Interaction with PSGL1/SELPLG is required for promoting recruitment and rolling of leukocytes. This interaction is dependent on the sialyl Lewis X glycan and tyrosine sulfation modifications of PSGL1. Sulfation on Y-51 of PSGL1 is



important for L-selectin binding.,tissue specificity:Expressed in B cell lines and T lymphocytes.,

Background

This gene encodes a cell surface adhesion molecule that belongs to a family of adhesion/homing receptors. The encoded protein contains a C-type lectin-like domain, a calcium-binding epidermal growth factor-like domain, and two short complement-like repeats. The gene product is required for binding and subsequent rolling of leucocytes on endothelial cells, facilitating their migration into secondary lymphoid organs and inflammation sites. Single-nucleotide polymorphisms in this gene have been associated with various diseases including immunoglobulin A nephropathy. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Oct 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using L-Selectin Monoclonal Antibody