



PIP4K2A Mouse mAb

Catalog No	YP-mAb-18961
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PIP4K2A PIP5K2 PIP5K2A
Immunogen	Synthesized peptide derived from human PIP4K2A
Specificity	This antibody detects endogenous levels of PIP4K2A at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal,Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PIPK;PI5P4KA;PIP5K2A;PIP5KIIA;PIP5KII-alpha;PIP4K2A;PIP4K2A PIP5K2 PIP5K2A;Phosphatidylinositol 5-phosphate 4-kinase type-2 alpha (1-phosphatidylinositol 5-phosphate 4-kinase 2-alpha) (Diphosphoinositide kinase 2-alpha) (PIP5KIII) (Phosphatidylinositol 5-phosphate 4-kinase type II alpha) (PI(5)P 4-kinase type II alpha) (PIP4KII-alpha) (PtdIns(4)P-5-kinase B isoform) (PtdIns(4)P-5-kinase C isoform) (PtdIns(5)P-4-kinase isoform 2-alpha)
Observed Band	46kDa
Calculated Molecular Weight	46kDa
Cell Pathway	Expressed ubiquitously, with high levels in the brain. Present in most tissues, except notably skeletal muscle and small intestine.
Tissue Specificity	
Function	Cell membrane . Nucleus . Lysosome . Cytoplasm . Photoreceptor inner segment . Cell projection, cilium, photoreceptor outer segment . May translocate from the cytosol to the cell membrane upon activation of tyrosine phosphorylation. May translocate from the inner to the outer segments of the rod photoreceptor cells in response to light (By similarity). Localization to the nucleus is modulated by the interaction with PIP4K2B. .

**Background**

Catalyzes the phosphorylation of phosphatidylinositol 5-phosphate (PtdIns5P) on the fourth hydroxyl of the myo-inositol ring, to form phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P₂) . Has both ATP- and GTP-dependent kinase activities . May exert its function by regulating the levels of PtdIns5P, which functions in the cytosol by increasing AKT activity and in the nucleus signals through ING2 . May regulate the pool of cytosolic PtdIns5P in response to the activation of tyrosine phosphorylation (By similarity). Required for lysosome-peroxisome membrane contacts and intracellular cholesterol transport through modulating peroxisomal PtdIns(4,5)P₂ level . In collaboration with PIP4K2B, has a role in mediating autophagy in times of nutrient stress (By similarity). Required for autophagosome-lysosome fusion and the regulation of cellular lipid metabolism . May be involved in thrombopoiesis, and the terminal maturation of megakaryocytes and regulation of their size (By similarity). Negatively regulates insulin signaling through a catalytic-independent mechanism . PIP4Ks interact with PIP5Ks and suppress PIP5K-mediated PtdIns(4,5)P₂ synthesis and insulin-dependent conversion to PtdIns(3,4,5)P₃ .

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images