



MUC1-C Mouse mAb

Catalog No	YP-mAb-19347
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	
Immunogen	Synthetic peptide. This information is considered to be commercially sensitive.
Specificity	
Formulation	
Source	Monoclonal,Mouse,IgG
Purification	Affinity purification
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NT5M DNT2;5'(3')-deoxyribonucleotidase;mitochondrial (5':3'-nucleotidase; mitochondrial) (Deoxy-5'-nucleotidase 2) (dNT-2);EMA;MCD;PEM;PUM;KL-6;MAM6;MCKD;PEMT;CD227;H23AG;MCKD1;MUC-1;ADMCKD;ADTKD2;Ca15-3;ADMCKD1;CA 15-3;MUC-1/X;MUC1/ZD;MUC-1/SEC;MUC1
Observed Band	17-25kDa
Calculated Molecular Weight	25kD
Cell Pathway	
Tissue Specificity	
Function	
Background	This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is



involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in multiple transcript variants.


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of various lysates using MUC1-C Mouse mAb (A19081) at 1:1000 dilution incubated at room temperature for 1.5 hours. Secondary antibody: HRP-conjugated Goat anti-Mouse IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Negative control (NC): HCT 116. Exposure time: 45 s