



eIF2 β Monoclonal Antibody

Catalog No	YP-mAb-03840
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	EIF2S2
Immunogen	Synthesized peptide derived from eIF2 β . at AA range: 10-90
Specificity	eIF2 β Monoclonal Antibody detects endogenous levels of eIF2 β protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	eIF2 β ;EIF2S2;Eukaryotic translation initiation factor 2 subunit 2;EIF2B;Eukaryotic translation initiation factor 2 subunit beta;eIF-2-beta
Observed Band	38kD
Calculated Molecular Weight	38kD
Cell Pathway	nucleus,cytoplasm,cytosol,eukaryotic translation initiation factor 2 complex,
Tissue Specificity	Amygdala,Brain,Epithelium,Lung,Muscle,Placenta,Pool
Function	function:eIF-2 functions in the early steps of protein synthesis by forming a ternary complex with GTP and initiator tRNA. This complex binds to a 40S ribosomal subunit, followed by mRNA binding to form a 43S preinitiation complex. Junction of the 60S ribosomal subunit to form the 80S initiation complex is preceded by hydrolysis of the GTP bound to eIF-2 and release of an eIF-2-GDP binary complex. In order for eIF-2 to recycle and catalyze another round of initiation, the GDP bound to eIF-2 must exchange with GTP by way of a reaction catalyzed by eIF-2B.,similarity:Belongs to the eIF-2-beta/eIF-5 family., subunit:Heterotrimer composed of an alpha, a beta and a gamma chain. Component of an EIF2 complex at least composed of CUGBP1, CALR, CALR3, EIF2S1, EIF2S2, HSP90B1 and HSPA5.,



Background

eukaryotic translation initiation factor 2 subunit beta(EIF2S2) Homo sapiens
Eukaryotic translation initiation factor 2 (EIF-2) functions in the early steps of protein synthesis by forming a ternary complex with GTP and initiator tRNA and binding to a 40S ribosomal subunit. EIF-2 is composed of three subunits, alpha, beta, and gamma, with the protein encoded by this gene representing the beta subunit. The beta subunit catalyzes the exchange of GDP for GTP, which recycles the EIF-2 complex for another round of initiation. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using eIF2 β Monoclonal Antibody