



Shh Monoclonal Antibody

Catalog No	YP-mAb-10842
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SHH
Immunogen	Synthesized peptide derived from human Shh
Specificity	This antibody detects endogenous levels of human Shh
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EC:3.1.1.-.;HHG 1;HHG1;HHG-1;HLP3;Shh;Sonic hedgehog protein (SHH;HHG-1) [Cleaved into: Sonic hedgehog protein N-product;Sonic hedgehog protein C-product]
Observed Band	40kD
Calculated Molecular Weight	40kD
Cell Pathway	Endoplasmic reticulum membrane . Golgi apparatus membrane . Co-localizes with HHAT in the ER and Golgi membrane. . ; [Sonic hedgehog protein N-product]: Cell membrane ; Lipid-anchor . The dual-lipidated sonic hedgehog protein N-product (ShhNp) is firmly tethered to the cell membrane where it forms multimers (PubMed:24522195). Further solubilization and release from the cell surface seem to be achieved through different mechanisms, including the interaction with DISP1 and SCUBE2, movement by lipoprotein particles, transport by cellular extensions called cytonemes or by the proteolytic removal of both terminal lipidated peptides (PubMed:26875496, PubMed:24522195). .
Tissue Specificity	Fetal lung,Plasma,
Function	disease:Defects in SHH are a cause of solitary median maxillary central incisor (SMMCI) [MIM:147250]. SMMCI is a rare dental anomaly characterized by the congenital absence of one maxillary central incisor.,disease:Defects in SHH are



the cause of holoprosencephaly type 3 (HPE3) [MIM:142945]. Holoprosencephaly (HPE) [MIM:236100] is the most common structural anomaly of the brain, in which the developing forebrain fails to correctly separate into right and left hemispheres. Holoprosencephaly is genetically heterogeneous and associated with several distinct facies and phenotypic variability. The majority of HPE3 cases are apparently sporadic, although clear examples of autosomal dominant inheritance have been described. Interestingly, up to 30% of obligate carriers of HPE3 gene in autosomal dominant pedigrees are clinically unaffected. disease: Defects in SHH are the cause of microphthalmia

Background

This gene encodes a protein that is instrumental in patterning the early embryo. It has been implicated as the key inductive signal in patterning of the ventral neural tube, the anterior-posterior limb axis, and the ventral somites. Of three human proteins showing sequence and functional similarity to the sonic hedgehog protein of *Drosophila*, this protein is the most similar. The protein is made as a precursor that is autocatalytically cleaved; the N-terminal portion is soluble and contains the signalling activity while the C-terminal portion is involved in precursor processing. More importantly, the C-terminal product covalently attaches a cholesterol moiety to the N-terminal product, restricting the N-terminal product to the cell surface and preventing it from freely diffusing throughout the developing embryo. Defects in this protein or in its signalling pathway are a cause of holoprosencephaly (HPE), a d

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using Shh Monoclonal Antibody