



HCFC2 mouse mAb

Catalog No	YP-mAb-11090
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	HCFC2
Immunogen	Synthesized peptide derived from human HCFC2 AA range: 347-397
Specificity	This antibody detects endogenous levels of HCFC2 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	C2 factor;HCF 2;HCF2;HCFC2;Host cell factor 2;host cell factor C2
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Cytoplasm . Nucleus .
Tissue Specificity	Highly expressed in testis. Detected at lower levels in spleen, thymus, prostate, ovary, small intestine and colon.
Function	similarity:Contains 3 fibronectin type-III domains.,similarity:Contains 4 Kelch repeats.,subunit:Binds MLL. Component of the MLL complex, at least composed of MLL, ASH2L, RBBP5, DPY30, WDR5, MEN1, HCFC1 and HCFC2.,tissue specificity:Highly expressed in testis. Detected at lower levels in spleen, thymus, prostate, ovary, small intestine and colon.,
Background	This gene encodes one of two proteins which interact with VP16, a herpes simplex virus protein that initiates virus infection. Both the encoded protein and the original Herpes host cell factor interact with VP16 through a beta-propeller domain. The original Herpes host cell factor, however, is effective at initiating viral infection while the encoded protein is not. Transcripts of varying length due to alternative polyadenylation signals have been described. [provided by RefSeq,




Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using HCFC2 mouse mAb