



PPIL1 mouse mAb

Catalog No	YP-mAb-11659
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PPIL1 CYPL1 CGI-124 UNQ2425/PRO4984
Immunogen	Synthesized peptide derived from human PPIL1 AA range: 73-123
Specificity	This antibody detects endogenous levels of PPIL1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CGI 124;CGI-124;CYPL1;EC:5.2.1.8;hCyPX;PPIL1;PPIL1 CYPL1 CGI-124 UNQ2425/PRO4984
Observed Band	
Calculated Molecular Weight	18kD
Cell Pathway	Nucleus .
Tissue Specificity	Ubiquitous, with the most abundant expression in heart and skeletal muscle.
Function	catalytic activity:Peptidylproline (omega=180) = peptidylproline (omega=0)., enzyme regulation:Inhibited by Cyclosporin A.,function:PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. May be involved in pre-mRNA splicing., similarity:Belongs to the cyclophilin-type PPlase family. PPIL1 subfamily., similarity:Contains 1 PPlase cyclophilin-type domain.,subunit:Identified in the spliceosome C complex, at least composed of AQR, ASCC3L1, C19orf29, CDC40, CDC5L, CRNKL1, DDX23, DDX41, DDX48, DDX5, DGCR14, DHX35, DHX38, DHX8, EFTUD2, FRG1, GPATC1, HNRPA1, HNRPA2B1, HNRPA3, HNRPC, HNRPF, HNRPH1, HNRPK, HNRPM, HNRPR, HNRPU, KIAA1160, KIAA1604, LSM2, LSM3, MAGOH, MORG1, MABPC1, PLRG1, PNN, PPIE, PPIL1, PPIL3, PPWD1, PRPF19, PRPF4B, PRPF6, PRPF8, RALY, RBM22,



RBM8A, RBMX, SART1, SF3A1, SF3A2, SF3A3, SF3B1, SF3B2, SF3B3,

Background

This gene is a member of the cyclophilin family of peptidylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PPIL1 mouse mAb