



Neuro D Monoclonal Antibody

Catalog No	YP-mAb-12755
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	NEUROD1
Immunogen	The antiserum was produced against synthesized peptide derived from human NEUROD1. AA range:240-289
Specificity	Neuro D Monoclonal Antibody detects endogenous levels of Neuro D protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NEUROD;Class A basic helix-loop-helix protein 3;bHLHa3;Neuro D;NEUROD1; Neurogenic differentiation factor 1
Observed Band	36kD
Calculated Molecular Weight	36kD
Cell Pathway	Cytoplasm . Nucleus . In pancreatic islet cells, shuttles to the nucleus in response to glucose stimulation (By similarity). Colocalizes with NR0B2 in the nucleus. .
Tissue Specificity	Eye,Retina,Rhabdomyosarcoma,
Function	disease:Defects in NEUROD1 are the cause of maturity onset diabetes of the young type 6 (MODY6) [MIM:606394]. MODY [MIM:606391] is characterized by an autosomal dominant mode of inheritance, onset during young adulthood and a primary defect in insulin secretion.,function:Differentiation factor required for dendrite morphogenesis and maintenance in the cerebellar cortex. Transcriptional activator. Binds to the insulin gene E-box.,PTM:Phosphorylated. In islet cells, phosphorylated on Ser-274 upon glucose stimulation; which may be required for nuclear localization. In activated neurons, phosphorylated on Ser-335; which promotes dendritic growth.,similarity:Contains 1 basic helix-loop-helix



(bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Heterodimer with TCF3/E47. Interacts with RREB1.,

Background

This gene encodes a member of the NeuroD family of basic helix-loop-helix (bHLH) transcription factors. The protein forms heterodimers with other bHLH proteins and activates transcription of genes that contain a specific DNA sequence known as the E-box. It regulates expression of the insulin gene, and mutations in this gene result in type II diabetes mellitus. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Neuro D Monoclonal Antibody