



ACTR-IIA Monoclonal Antibody

Catalog No	YP-mAb-13135
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ACVR2A
Immunogen	The antiserum was produced against synthesized peptide derived from human ACTR-IIA. AA range:10-59
Specificity	ACTR-IIA Monoclonal Antibody detects endogenous levels of ACTR-IIA protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ACTR-IIA;ACVR2A;Activin receptor type-2A;ACVR2;Activin receptor type IIA; ACTRIIA
Observed Band	48kD
Calculated Molecular Weight	48kD
Cell Pathway	Membrane; Single-pass type I membrane protein.
Tissue Specificity	Mammary gland,Testis,
Function	catalytic activity:ATP + [receptor-protein] = ADP + [receptor-protein] phosphate., cofactor:Magnesium or manganese.,function:On ligand binding, forms a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. Receptor for activin A, activin B and inhibin A.,similarity:Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. TGFB receptor subfamily., similarity:Contains 1 protein kinase domain.,subunit:Interacts with AIP1. Part of a complex consisting of AIP1, ACVR2A, ACVR1B and SMAD3.,



Background

This gene encodes a receptor that mediates the functions of activins, which are members of the transforming growth factor-beta (TGF-beta) superfamily involved in diverse biological processes. The encoded protein is a transmembrane serine-threonine kinase receptor which mediates signaling by forming heterodimeric complexes with various combinations of type I and type II receptors and ligands in a cell-specific manner. The encoded type II receptor is primarily involved in ligand-binding and includes an extracellular ligand-binding domain, a transmembrane domain and a cytoplasmic serine-threonine kinase domain. This gene may be associated with susceptibility to preeclampsia, a pregnancy-related disease which can result in maternal and fetal morbidity and mortality. Alternative splicing results in multiple transcript variants of this gene. [provided by RefSeq, Jun 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using ACTR-IIA Monoclonal Antibody