



CXCR-7 Monoclonal Antibody

Catalog No	YP-mAb-13189
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CXCR7
Immunogen	The antiserum was produced against synthesized peptide derived from human CXCR7. AA range:311-360
Specificity	CXCR-7 Monoclonal Antibody detects endogenous levels of CXCR-7 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	RDC1;CXCR7;RDC-1;CMKOR1;CXC-R7;CXCR-7;GPR159;ACKR3;C-X-C chemokine receptor type 7;Chemokine orphan receptor 1;G-protein coupled receptor 159;G-protein coupled receptor RDC1 homolog
Observed Band	41kD
Calculated Molecular Weight	41kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein . Early endosome . Recycling endosome . Predominantly localizes to endocytic vesicles, and upon stimulation by the ligand is internalized via clathrin-coated pits in a beta-arrestin-dependent manner. Once internalized, the ligand dissociates from the receptor, and is targeted to degradation while the receptor is recycled back to the cell membrane. .
Tissue Specificity	Expressed in monocytes, basophils, B-cells, umbilical vein endothelial cells (HUVEC) and B-lymphoblastoid cells. Lower expression detected in CD4+ T-lymphocytes and natural killer cells. In the brain, detected in endothelial cells and capillaries, and in mature neurons of the frontal cortex and hippocampus. Expressed in tubular formation in the kidney. Highly expressed in astroglial tumor endothelial, microglial and glioma cells. Expressed at low levels in normal CD34+ progenitor cells, but at very high levels in several myeloid malignant cell lines. Expressed in breast carcinomas but not in normal breast tissue (at protein



level).

Function

caution:Was originally (PubMed:1675791) thought to be the receptor for VIP.,
function:Receptor for CXCL12/SDF1. Acts as coreceptor with CXCR4 for a
restricted number of HIV isolates.,online information:CXC chemokine receptors
entry,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue
specificity:Expressed in monocytes, basophils, and B-cells. Lower expression in
CD4+ T-lymphocytes and natural killer cells.,

Background

This gene encodes a member of the G-protein coupled receptor family. Although
this protein was earlier thought to be a receptor for vasoactive intestinal peptide
(VIP), it is now considered to be an orphan receptor, in that its endogenous
ligand has not been identified. The protein is also a coreceptor for human
immunodeficiency viruses (HIV). Translocations involving this gene and HMGA2
on chromosome 12 have been observed in lipomas. [provided by RefSeq, Jul
2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For
more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CXCR-7
Monoclonal Antibody