



MerTK/Tyro3 Monoclonal Antibody

Catalog No	YP-mAb-13415
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	MERTK/TYRO3
Immunogen	The antiserum was produced against synthesized peptide derived from human MER/SKY. AA range:716-765
Specificity	MerTK/Tyro3 Monoclonal Antibody detects endogenous levels of MerTK/Tyro3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	c mer;EC:2.7.10.1;MER;Proto oncogene c Mer;Proto-oncogene c-Mer;MerTK/Tyro3;Tyrosine-protein kinase Mer/Tyrosine-protein kinase receptor TYRO3;MERTK;Tyrosine-protein kinase Mer;Receptor tyrosine kinase MerTK;TYRO3;BYK;DTK;RSE;SKY;Tyrosine-protein kinase receptor TYRO3;Tyrosine-protein kinase DTK;Tyrosine-protein kinase RSE;Tyrosine-protein kin
Observed Band	97kD
Calculated Molecular Weight	97kD
Cell Pathway	Cell membrane ; Single-pass type I membrane protein .
Tissue Specificity	Not expressed in normal B- and T-lymphocytes but is expressed in numerous neoplastic B- and T-cell lines. Highly expressed in testis, ovary, prostate, lung, and kidney, with lower expression in spleen, small intestine, colon, and liver.
Function	catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,disease:Defects in MERTK are a cause of retinitis pigmentosa (RP) [MIM:268000]. RP that leads to degeneration of retinal photoreceptor cells. Patients typically have night vision blindness and loss of midperipheral visual field. As their condition progresses, they lose their far peripheral visual field and eventually central vision as well.,function:In case of filovirus infection, seems to



function as a cell entry factor.,online information:Retina International's Scientific Newsletter,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. AXL/UFO subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 fibronectin type-III domains.,similarity:Contains 2 Ig-like C2-type (imm

Background

This gene is a member of the MER/AXL/TYRO3 receptor kinase family and encodes a transmembrane protein with two fibronectin type-III domains, two Ig-like C2-type (immunoglobulin-like) domains, and one tyrosine kinase domain. Mutations in this gene have been associated with disruption of the retinal pigment epithelium (RPE) phagocytosis pathway and onset of autosomal recessive retinitis pigmentosa (RP). [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using MerTK/Tyro3 Monoclonal Antibody