



Relaxin Receptor 3 Monoclonal Antibody

Catalog No	YP-mAb-13668
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	RXFP3
Immunogen	The antiserum was produced against synthesized peptide derived from human RXFP3. AA range:161-210
Specificity	Relaxin Receptor 3 Monoclonal Antibody detects endogenous levels of Relaxin Receptor 3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	G protein-coupled receptor SALPR;GPCR135;G-protein coupled receptor GPCR135;Relaxin 3 receptor 1;Relaxin-3 receptor 1;Relaxin Receptor 3;RXFP3;RLN3R1;SALPR;RLN3 receptor 1;Relaxin family peptide receptor 3;Somatostatin- and angiotensin-like peptide receptor
Observed Band	51kD
Calculated Molecular Weight	51kD
Cell Pathway	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Expressed predominantly in brain regions. Highest expression in substantia nigra and pituitary, followed by hippocampus, spinal cord, amygdala, caudate nucleus and corpus callosum, quite low level in cerebellum. In peripheral tissues, relatively high levels in adrenal glands, low levels in pancreas, salivary gland, placenta, mammary gland and testis.
Function	function:Receptor for relaxin-3. Binding of the ligand inhibit cAMP accumulation., similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed predominantly in brain regions. Highest expression in substantia nigra and pituitary, followed by hippocampus, spinal cord, amygdala, caudate nucleus and corpus callosum, quite low level in cerebellum. In



peripheral tissues, relatively high levels in adrenal glands, low levels in pancreas, salivary gland, placenta, mammary gland and testis.,

Background

function: Receptor for relaxin-3. Binding of the ligand inhibit cAMP accumulation., similarity: Belongs to the G-protein coupled receptor 1 family., tissue specificity: Expressed predominantly in brain regions. Highest expression in substantia nigra and pituitary, followed by hippocampus, spinal cord, amygdala, caudate nucleus and corpus callosum, quite low level in cerebellum. In peripheral tissues, relatively high levels in adrenal glands, low levels in pancreas, salivary gland, placenta, mammary gland and testis.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

