



IL-3R α Monoclonal Antibody

Catalog No	YP-mAb-13742
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	IL3RA
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human IL3RA. AA range:241-290
Specificity	IL-3R α Monoclonal Antibody detects endogenous levels of IL-3R α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	IL-3R α ;IL3RA;Interleukin-3 receptor subunit alpha;IL3R;IL-3 receptor subunit alpha;IL-3R subunit alpha;IL-3R-alpha;IL-3RA;CD123
Observed Band	42kD
Calculated Molecular Weight	42kD
Cell Pathway	Membrane; Single-pass type I membrane protein.
Tissue Specificity	Brain,Lung,
Function	domain:The box 1 motif is required for JAK interaction and/or activation., domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding., function:This is a receptor for interleukin-3.,miscellaneous:The gene encoding for this protein is located in the pseudoautosomal region 1 (PAR1) of X and Y chromosomes.,similarity:Belongs to the type I cytokine receptor family. Type 5 subfamily.,subunit:Heterodimer of an alpha and a beta subunit. The beta subunit is common to the IL3, IL5 and GM-CSF receptors.,



Background

The protein encoded by this gene is an interleukin 3 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL3 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL3. This gene and the gene encoding the colony stimulating factor 2 receptor alpha chain (CSF2RA) form a cytokine receptor gene cluster in a X-Y pseudoautosomal region on chromosomes X or Y. Alternatively spliced transcript variants encoding distinct isoforms have been found. [provided by RefSeq, Jun 2012],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using IL-3Ra Monoclonal Antibody