



HLA-DOβ Monoclonal Antibody

Catalog No	YP-mAb-13930
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	HLA-DOB
Immunogen	The antiserum was produced against synthesized peptide derived from human HLA-DOB. AA range:1-50
Specificity	HLA-DOβ Monoclonal Antibody detects endogenous levels of HLA-DOβ protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HLA-B;HLA B;HLA class I histocompatibility antigen;B alpha chain;Human leukocyte antigen B;HLA-DOβ;HLA-DOB;HLA class II histocompatibility antigen DO beta chain;HLA class II histocompatibility antigen;DO beta chain;MHC class II antigen DOB
Observed Band	30kD
Calculated Molecular Weight	30kD
Cell Pathway	Endosome membrane; Single-pass type I membrane protein. Lysosome membrane; Single-pass type I membrane protein. Complexes with HLA-DM molecule during intracellular transport and in endosomal/lysosomal compartments. Heterotetramerization is necessary to exit the ER.
Tissue Specificity	B-cell,
Function	function:Important modulator in the HLA class II restricted antigen presentation pathway by interaction with the HLA-DM molecule.,polymorphism:The following alleles of DOB are known: DOB*0101, DOB*0102, DOB*0103 and DOB*0104. The sequence shown is that of DOB*0101.,similarity:Belongs to the MHC class II family.,similarity:Contains 1 Ig-like C1-type (immunoglobulin-like) domain., subunit:Heterodimer of an alpha chain (DOA) and a beta chain (DOB).,



Background

HLA-DOB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DOA) and a beta chain (DOB), both anchored in the membrane. It is located in intracellular vesicles. DO suppresses peptide loading of MHC class II molecules by inhibiting HLA-DM. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using HLA-DO β Monoclonal Antibody