



CD85d Monoclonal Antibody

Catalog No	YP-mAb-14058
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	LILRB2
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human LILRB2. AA range:121-170
Specificity	CD85d Monoclonal Antibody detects endogenous levels of CD85d protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	LILRB2;CD85 antigen-like family member D;CD85d;ILT 4;ILT4;Leukocyte immunoglobulin-like receptor subfamily B member 2;LIR2;MIR10;LIR-2; Leukocyte immunoglobulin-like receptor 2;Immunoglobulin-like transcript 4;ILT-4; Monocyte/macrophage immunoglobulin-like receptor 10;MIR-10
Observed Band	66kD
Calculated Molecular Weight	66kD
Cell Pathway	Cell membrane ; Single-pass type I membrane protein.
Tissue Specificity	Expressed in monocytes and at lower levels in myeloid and plasmacytoid dendritic cells. Expressed in tolerogenic IL10-producing dendritic cells (PubMed:20448110). Expressed in myeloid-derived suppressor cells during pregnancy (PubMed:27859042). Detected at low levels in natural killer (NK) cells. Expressed in B cells.
Function	domain:Contains 3 copies of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.,function:Receptor for class I MHC antigens. Recognizes a broad spectrum of HLA-A, HLA-B, HLA-C



and HLA-G alleles. Involved in the down-regulation of the immune response and the development of tolerance. Competes with CD8A for binding to class I MHC antigens. Inhibits FCGR1A-mediated phosphorylation of cellular proteins and mobilization of intracellular calcium ions. PTM: Phosphorylated on tyrosine residues. Dephosphorylated by PTPN6. similarity: Contains 4 Ig-like C2-type (immunoglobulin-like) domains. subunit: Binds PTPN6 when phosphorylated. Binds FCGR1A. tissue specificity: Expressed on monocytes and B-cells, a

Background

This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CD85d Monoclonal Antibody