



CD93 Monoclonal Antibody

Catalog No	YP-mAb-14060
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CD93
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human CD93. AA range:191-240
Specificity	CD93 Monoclonal Antibody detects endogenous levels of CD93 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	C1QR1;Matrix-remodeling-associated protein 4;C1qRp;C1qR(p);C1qR;CD93; Complement component C1q receptor;MXRA4;C1q/MBL/SPA receptor;CDw93; Complement component 1 q subcomponent receptor 1
Observed Band	110-120kD
Calculated Molecular Weight	110-120kD
Cell Pathway	Membrane; Single-pass type I membrane protein.
Tissue Specificity	Highly expressed in endothelial cells, platelets, cells of myeloid origin, such as monocytes and neutrophils. Not expressed in cells of lymphoid origin.
Function	caution:Has been sometimes referred to as a collectin receptor., caution:PubMed:11994479 reported that C1q is not a ligand for C1QR1., function:Receptor (or element of a larger receptor complex) for C1q, mannose-binding lectin (MBL2) and pulmonary surfactant protein A (SPA). May mediate the enhancement of phagocytosis in monocytes and macrophages upon interaction with soluble defense collagens. May play a role in intercellular adhesion.,PTM:N- and O-glycosylated.,similarity:Contains 1 C-type lectin domain.,similarity:Contains 5 EGF-like domains.,subunit:Interacts with HCV core protein.,tissue specificity:Highly expressed in endothelial cells, platelets, cells of



myeloid origin, such as monocytes and neutrophils. Not expressed in cells of lymphoid origin.,

Background

The protein encoded by this gene is a cell-surface glycoprotein and type I membrane protein that was originally identified as a myeloid cell-specific marker. The encoded protein was once thought to be a receptor for C1q, but now is thought to instead be involved in intercellular adhesion and in the clearance of apoptotic cells. The intracellular cytoplasmic tail of this protein has been found to interact with moesin, a protein known to play a role in linking transmembrane proteins to the cytoskeleton and in the remodelling of the cytoskeleton. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CD93 Monoclonal Antibody