



CD209 Monoclonal Antibody

Catalog No	YP-mAb-14077
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CD209
Immunogen	Synthesized peptide derived from CD209 antigen at AA range: 261-310
Specificity	CD209 Monoclonal Antibody detects endogenous levels of CD209 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DC-SIGN;CLEC4L;C-type lectin domain family 4 member L;DC SIGN;DC; CD209;CD209 antigen;Dendritic cell-specific ICAM-3-grabbing non-integrin 1; DC-SIGN1
Observed Band	45kD
Calculated Molecular Weight	45kD
Cell Pathway	[Isoform 1]: Cell membrane ; Single-pass type II membrane protein .; [Isoform 2]: Cell membrane ; Single-pass type II membrane protein .; [Isoform 3]: Cell membrane ; Single-pass type II membrane protein .; [Isoform 4]: Cell membrane ; Single-pass type II membrane protein .; [Isoform 5]: Cell membrane ; Single-pass type II membrane protein .; [Isoform 6]: Secreted .; [Isoform 7]: Secreted .; [Isoform 8]: Secreted .; [Isoform 9]: Secreted .; [Isoform 10]: Secreted .; [Isoform 11]: Secreted .; [Isoform 12]: Secreted .
Tissue Specificity	Predominantly expressed in dendritic cells and in DC-residing tissues. Also found in placental macrophages, endothelial cells of placental vascular channels, peripheral blood mononuclear cells, and THP-1 monocytes.
Function	alternative products:Additional isoforms seem to exist. Several splicing events may be used independently in a modular way. Deletion of the transmembrane domain encoding exon through alternative splicing produces soluble isoforms,



caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,domain:The tandem repeat domain, also called neck domain, mediates oligomerization.,function:On DCs it is a high affinity receptor for ICAM2 and ICAM3 by binding to mannose-like carbohydrates. May act as a DC rolling receptor that mediates transendothelial migration of DC presursors from blood to tissues by binding endothelial ICAM2. Seems to regulate DC-induced T-cell proliferation by binding to ICAM3 on T-cells in the immunological synapse formed between DC and T-cells.,function:Pathogen-recognition receptor expressed on the sur

Background

This gene encodes a transmembrane receptor and is often referred to as DC-SIGN because of its expression on the surface of dendritic cells and macrophages. The encoded protein is involved in the innate immune system and recognizes numerous evolutionarily divergent pathogens ranging from parasites to viruses with a large impact on public health. The protein is organized into three distinct domains: an N-terminal transmembrane domain, a tandem-repeat neck domain and C-type lectin carbohydrate recognition domain. The extracellular region consisting of the C-type lectin and neck domains has a dual function as a pathogen recognition receptor and a cell adhesion receptor by binding carbohydrate ligands on the surface of microbes and endogenous cells. The neck region is important for homo-oligomerization which allows the receptor to bind multivalent ligands with high avidity. Variations in the number of 23 amino a


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CD209 Monoclonal Antibody