



DGK-α Monoclonal Antibody

Catalog No	YP-mAb-14721
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	DGKA
Immunogen	The antiserum was produced against synthesized peptide derived from human DGKA. AA range:304-353
Specificity	DGK-α Monoclonal Antibody detects endogenous levels of DGK-α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DAGK;DAGK1;DGK alpha;DGK-alpha;EC:2.7.1.107;DGK-α;DGKA; Diacylglycerol kinase alpha;DAG kinase alpha;80 kDa diacylglycerol kinase; Diglyceride kinase alpha
Observed Band	80kD
Calculated Molecular Weight	80kD
Cell Pathway	Cytoplasm, cytosol .
Tissue Specificity	Expressed in lymphocytes.
Function	catalytic activity:ATP + 1,2-diacylglycerol = ADP + 1,2-diacyl-sn-glycerol 3-phosphate.,enzyme regulation:Stimulated by calcium and phosphatidylserine. Phosphorylated by protein kinase C.,function:Upon cell stimulation converts the second messenger diacylglycerol into phosphatidate, initiating the resynthesis of phosphatidylinositols and attenuating protein kinase C activity.,similarity:Belongs to the eukaryotic diacylglycerol kinase family.,similarity:Contains 1 DAGKc domain.,similarity:Contains 2 EF-hand domains.,similarity:Contains 2 phorbol-ester/DAG-type zinc fingers.,subunit:Monomer.,tissue specificity:Lymphocytes and oligodendroglial cells.,



Background

The protein encoded by this gene belongs to the eukaryotic diacylglycerol kinase family. It acts as a modulator that competes with protein kinase C for the second messenger diacylglycerol in intracellular signaling pathways. It also plays an important role in the resynthesis of phosphatidylinositols and phosphorylating diacylglycerol to phosphatidic acid. Alternative splicing occurs at this locus and four transcript variants encoding the same protein have been identified. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using DGK- α Monoclonal Antibody