




IKK α / β Monoclonal Antibody

Catalog No	YP-mAb-14781
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CHUK/IKBKB
Immunogen	The antiserum was produced against synthesized peptide derived from human IKK-alpha/beta. AA range:141-190
Specificity	IKK α / β Monoclonal Antibody detects endogenous levels of IKK α / β protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	chuk;CHUK1;Conserved Helix Loop Helix Ubiquitous Kinase;Conserved helix loop ubiquitous kinase;Conserved helix-loop-helix ubiquitous kinase;I Kappa B Kinase 1;I Kappa B Kinase Alpha;l-kappa-B kinase 1;l-kappa-B kinase alpha;l-kappaB kinase;lB kinase alpha subunit;lBKA;IKK 1;IKK A;IKK a kinase;IKK-A;IKK-alpha;IKK1;IKKA;IKKA_HUMAN;Inhibitor Of Kappa Light Polypeptide Gene Enhancer In B Cells;Inhibitor Of Nuclear Factor Kappa B Kinase Alpha Subunit;Inhibitor of nuclear factor kappa-B kinase subunit alpha;NFKBIKA;Nuclear Factor Kappa B Inhibitor Kinase Alpha;Nuclear factor NF kappa B inhibitor kinase alpha;Nuclear factor NF-kappa-B inhibitor kinase alpha;Nuclear factor NFkappaB inhibitor kinase alpha;Nuclear Factor Of Kappa Light Chain Gene Enhancer In B Cells Inhibitor;TCF-16;TCF16;Transcription factor 16;l kappa B kinase 2;l kappa B kinase beta;l-kappa-B kinase 2;l-kappa-B-kinase beta;lBKB;IKK beta;IKK-B;IKK-beta;IKK2;IKKB;IKKB_HUMAN;IMD15;kinase beta;Inhibitor of nuclear factor kappa-B kinase subunit beta;NFKBIKB;Nuclear factor NF-kappa-B inhibitor kinase beta;IKK α / β ;CHUK/IKBKB;Nuclear factor NF-kappa-B
Observed Band	85kD
Calculated Molecular Weight	85kD
Cell Pathway	Cytoplasm . Nucleus . Shuttles between the cytoplasm and the nucleus.



Tissue Specificity	Widely expressed.
Function	catalytic activity:ATP + [I-kappa-B protein] = ADP + [I-kappa-B phosphoprotein]., enzyme regulation:Activated when phosphorylated and inactivated when dephosphorylated.,function:Acts as part of the IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor. As part of the non-canonical pathway of NF-kappa-B activation, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFKB2/p100 associated with RelB, inducing its proteolytic processing to NFKB2/p52 and the formation of NF-kappa-B RelB-p52 complexes. Also phosphorylates NCOA3. Phosphorylates 'Ser-10' of histone H3 at NF-kappa-B-regulated promoters during inflammatory responses triggered by cytokines.,PTM:Phosphorylated by MAP3K14/NIK, AKT and to a lesser extent by MEKK
Background	This gene encodes a member of the serine/threonine protein kinase family. The encoded protein, a component of a cytokine-activated protein complex that is an inhibitor of the essential transcription factor NF-kappa-B complex, phosphorylates sites that trigger the degradation of the inhibitor via the ubiquination pathway, thereby activating the transcription factor. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using IKK α / β Monoclonal Antibody