



NARF Monoclonal Antibody

Catalog No	YP-mAb-14872
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	NARF
Immunogen	The antiserum was produced against synthesized peptide derived from human NARF. AA range:262-311
Specificity	NARF Monoclonal Antibody detects endogenous levels of NARF protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	IOP2;NARF;Nuclear prelamin A recognition factor;Iron-only hydrogenase-like protein 2
Observed Band	51kD
Calculated Molecular Weight	51kD
Cell Pathway	Nucleus .
Tissue Specificity	Ubiquitous. Predominantly expressed in skeletal muscle, heart and brain.
Function	PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., similarity:Belongs to the NARF family.,subunit:Interacts with LMNA and binds to the farnesylated C-terminal domain.,tissue specificity:Ubiquitous. Predominantly expressed in skeletal muscle, heart and brain.,
Background	Several proteins have been found to be prenylated and methylated at their carboxyl-terminal ends. Prenylation was initially believed to be important only for membrane attachment. However, another role for prenylation appears to be its importance in protein-protein interactions. The only nuclear proteins known to be prenylated in mammalian cells are prelamin A- and B-type lamins. Prelamin A is farnesylated and carboxymethylated on the cysteine residue of a carboxyl-



terminal CaaX motif. This post-translationally modified cysteine residue is removed from prelamin A when it is endoproteolytically processed into mature lamin A. The protein encoded by this gene binds to the prenylated prelamin A carboxyl-terminal tail domain. It may be a component of a prelamin A endoprotease complex. The encoded protein is located in the nucleus, where it partially colocalizes with the nuclear lamina. It

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of various cells using NARF Monoclonal Antibody