



Wee 2 Monoclonal Antibody

Catalog No	YP-mAb-15033
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	WEE2
Immunogen	The antiserum was produced against synthesized peptide derived from human WEE2. AA range:151-200
Specificity	Wee 2 Monoclonal Antibody detects endogenous levels of Wee 2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	WEE1 homolog 2 (S. pombe);Wee1 like protein kinase 1B;Wee1 like protein kinase 2;WEE1B;Wee1B kinase;WEE2;Wee 2;Wee1-like protein kinase 2;Wee1-like protein kinase 1B
Observed Band	60kD
Calculated Molecular Weight	60kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in oocytes (at protein level) (PubMed:29606300). May also be expressed in testis (PubMed:11029659).
Function	catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,function:Phosphorylates and inhibits CDC2. May act as a negative regulator of entry into mitosis (G2 to M transition).,PTM:Phosphorylated ., similarity:Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. WEE1 subfamily.,similarity:Contains 1 protein kinase domain.,tissue specificity:Expressed in testis.,



Background

catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,function:Phosphorylates and inhibits CDC2. May act as a negative regulator of entry into mitosis (G2 to M transition).,PTM:Phosphorylated ., similarity:Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. WEE1 subfamily.,similarity:Contains 1 protein kinase domain.,tissue specificity:Expressed in testis.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Wee 2 Monoclonal Antibody