



Brn-3a Monoclonal Antibody

Catalog No	YP-mAb-15749
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	POU4F1
Immunogen	The antiserum was produced against synthesized peptide derived from human POU4F1. AA range:311-360
Specificity	Brn-3a Monoclonal Antibody detects endogenous levels of Brn-3a protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Brain specific homeobox/POU domain protein 3A;Brain-3A;Brain-specific homeobox/POU domain protein 3A;Brn 3.;BRN 3A;Brn-3A;Brn3;BRN3A;class 4; FLJ13449;Homeobox/POU domain protein RDC 1;Homeobox/POU domain protein RDC-1;Homeobox/POU domain protein RDC1;Oct T1;Oct-T1;OctT1; PO4F1_HUMAN;POU class 4 homeobox 1;POU domain;POU Domain Class 4 Transcription Factor 1;Pou4f1;RDC 1;RDC1;transcription factor 1; transcription factor 1
Observed Band	42kD
Calculated Molecular Weight	42kD
Cell Pathway	Nucleus . Cytoplasm .
Tissue Specificity	Expressed in the brain and the retina. Present in the developing brain, spinal cord and eye.
Function	developmental stage:Expression peaks early in embryogenesis (day 13.5) and is undetectable 14 days after birth.,function:Probable transcription factor which may play a role in the regulation of specific gene expression within a subset of neuronal lineages. May play a role in determining or maintaining the identities of a small subset of visual system neurons.,similarity:Belongs to the POU



transcription factor family. Class-4 subfamily.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 1 POU-specific domain.,tissue specificity:Brain. Seems to be specific to the retina. Present in the developing brain, spinal cord and eye.,

Background

This gene encodes a member of the POU-IV class of neural transcription factors. This protein is expressed in a subset of retinal ganglion cells and may be involved in the developing sensory nervous system. This protein may also promote the growth of cervical tumors. A translocation of this gene is associated with some adult acute myeloid leukemias. [provided by RefSeq, Mar 2012],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Brn-3a Monoclonal Antibody