



IGFBP3 Monoclonal Antibody

Catalog No	YP-mAb-15925
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	IGFBP3
Immunogen	The antiserum was produced against synthesized peptide derived from human IGFBP-3. AA range:151-200
Specificity	IGFBP3 Monoclonal Antibody detects endogenous levels of IGFBP3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	BP 53;IBP 3;IBP3;IBP-3;IGF binding protein 3;IGFBP3;Insulin-like growth factor-binding protein 3;IGF-binding protein 3;IGFBP-3
Observed Band	31kD
Calculated Molecular Weight	31kD
Cell Pathway	Secreted .
Tissue Specificity	Expressed by most tissues. Present in plasma.
Function	developmental stage:IGFBP3 levels are higher during extrauterine life and peak during puberty.,domain:The thyroglobulin type-1 domain mediates interaction with HN.,function:IGF-binding proteins prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors.,induction:IGFBP3 levels increase in the presence of IGF1, insulin and other growth-stimulating factors such as growth hormone, epidermal growth factor, and phorbol esters.,online information:The Singapore human mutation and polymorphism database,similarity:Contains 1 IGFBP N-terminal domain.,similarity:Contains 1 thyroglobulin type-1 domain.,subunit:Interacts with XLKD1 (By similarity). Binds IGF2 more than IGF1. Forms a ternary complex of about



140 to 150 kDa with IGF1 or IGF2 and a 85 k

Background

This gene is a member of the insulin-like growth factor binding protein (IGFBP) family and encodes a protein with an IGFBP domain and a thyroglobulin type-I domain. The protein forms a ternary complex with insulin-like growth factor acid-labile subunit (IGFALS) and either insulin-like growth factor (IGF) I or II. In this form, it circulates in the plasma, prolonging the half-life of IGFs and altering their interaction with cell surface receptors. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using IGFBP3 Monoclonal Antibody