



IL-20R α Monoclonal Antibody

Catalog No	YP-mAb-15934
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	IL20RA
Immunogen	The antiserum was produced against synthesized peptide derived from human IL-20R α . AA range:173-222
Specificity	IL-20R α Monoclonal Antibody detects endogenous levels of IL-20R α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	class II cytokine receptor ZCYTOR7;CRF2 8;CRF2-8;Cytokine receptor class II member 8;Cytokine receptor class-II member 8;Cytokine receptor family 2 member 8;I2RA HUMAN;IL 2R alpha;IL 2R1;IL-2 receptor subunit alpha;IL-2R-alpha;IL-2R1;IL-2RA;IL2RA;Interleukin 2 receptor alpha chain;Interleukin 2 receptor;alpha;interleukin-2 receptor I;Interleukin-2 receptor subunit alpha;ZcytoR7;IL-20R α ;IL20RA;Interleukin-20 receptor subunit alpha;IL-20 receptor subunit alpha;IL-20R-alpha;IL-20RA;IL-20R1
Observed Band	62kD
Calculated Molecular Weight	62kD
Cell Pathway	Membrane ; Single-pass type I membrane protein .
Tissue Specificity	Widely expressed with highest levels in skin and testis and high levels in brain. Highly expressed in psoriatic skin.
Function	function:The IL20RA/IL20RB dimer is a receptor for IL19, IL20 and IL24. The IL20RA/IL10RB dimer is a receptor for IL26.,similarity:Belongs to the type II cytokine receptor family.,similarity:Contains 2 fibronectin type-III domains., subunit:Heterodimer with IL20RB and heterodimer with IL10RB.,tissue specificity:Widely expressed with highest levels in skin and testis and high levels



in brain. Highly expressed in psoriatic skin.,

Background

This gene encodes a member of the type II cytokine receptor family. The encoded protein is a subunit of the receptor for interleukin 20, a cytokine that may be involved in epidermal function. The interleukin 20 receptor is a heterodimeric complex consisting of the encoded protein and interleukin 20 receptor beta. This gene and interleukin 20 receptor beta are highly expressed in skin, and are upregulated in psoriasis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using IL-20Rα Monoclonal Antibody