



VEGF-C Monoclonal Antibody

Catalog No	YP-mAb-16012
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	VEGFC
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human VEGFC. AA range:91-140
Specificity	VEGF-C Monoclonal Antibody detects endogenous levels of VEGF-C protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	VEGFC;vascular endothelial growth factor C;Vascular endothelial growth factor-related protein;VEGF C;VEGF-C;Flt4 ligand;Flt4-L;VRP
Observed Band	42kD
Calculated Molecular Weight	42kD
Cell Pathway	Secreted.
Tissue Specificity	Spleen, lymph node, thymus, appendix, bone marrow, heart, placenta, ovary, skeletal muscle, prostate, testis, colon and small intestine and fetal liver, lung and kidney, but not in peripheral blood lymphocyte.
Function	function:Growth factor active in angiogenesis, and endothelial cell growth, stimulating their proliferation and migration and also has effects on the permeability of blood vessels. May function in angiogenesis of the venous and lymphatic vascular systems during embryogenesis, and also in the maintenance of differentiated lymphatic endothelium in adults. Binds and activates VEGFR-2 (Flk1) and VEGFR-3 (Flt4) receptors.,PTM:Undergoes a complex proteolytic maturation which generates a variety of processed secreted forms with increased activity toward VEGFR-3, but only the fully processed form could activate VEGFR-2. VEGF-C first form an antiparallel homodimer linked by



disulfide bonds. Before secretion, a cleavage occurs between Arg-227 and Ser-228 producing an heterotetramer. The next extracellular step of the processing removes the N-terminal propeptide. Finally the mature VEGF-C is compos

Background

The protein encoded by this gene is a member of the platelet-derived growth factor/vascular endothelial growth factor (PDGF/VEGF) family. The encoded protein promotes angiogenesis and endothelial cell growth, and can also affect the permeability of blood vessels. The proprotein is further cleaved into a fully processed form that can bind and activate VEGFR-2 and VEGFR-3 receptors. [provided by RefSeq, Apr 2014],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

E



Western Blot analysis of various cells using VEGF-C Monoclonal Antibody