



ANG I Monoclonal Antibody

Catalog No	YP-mAb-16048
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ANG
Immunogen	The antiserum was produced against synthesized peptide derived from the N-terminal region of human N-ternal ANG. AA range:11-60
Specificity	ANG I Monoclonal Antibody detects endogenous levels of ANG I protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ALS9;AMYOTROPHIC LATERAL SCLEROSIS;ANG 1;ANG;ANG I;ANG1;ANGI;ANGI_HUMAN;Angiogenin;Angiogenin ribonuclease RNase A family;5;Epididymis luminal protein 168;HEL168;MGC22466;MGC71966;Ribonuclease 5;Ribonuclease RNase A Family 5;RNase 5;RNASE4;RNASE5;Ang-1
Observed Band	17kD
Calculated Molecular Weight	17kD
Cell Pathway	Cytoplasmic vesicle, secretory vesicle lumen . Secreted . Nucleus . Nucleus, nucleolus . Rapidly endocytosed by target cells and translocated to the nucleus where it accumulates in the nucleolus and binds to DNA (PubMed:12051708). .
Tissue Specificity	Expressed predominantly in the liver. Also detected in endothelial cells and spinal cord neurons.
Function	caution:It is uncertain whether Met-1 or Met-3 is the initiator.,developmental stage:Low level expression in the developing fetus, increased in the neonate, and maximal in the adult.,disease:Defects in ANG are the cause of susceptibility to amyotrophic lateral sclerosis type 9 (ALS9) [MIM:611895]. ALS is a degenerative disorder of motor neurons in the cortex, brain stem and spinal cord. ALS is characterized by muscular weakness and atrophy.,function:May function as a tRNA-specific ribonuclease that binds to actin on the surface of



endothelial cells; once bound, angiogenin is endocytosed and translocated to the nucleus, thereby promoting the endothelial invasiveness necessary for blood vessel formation. Angiogenin induces vascularization of normal and malignant tissues. Abolishes protein synthesis by specifically hydrolyzing cellular tRNAs., online information:Angiogenin,similarity:Belongs

Background

angiogenin(ANG) Homo sapiens The protein encoded by this gene is an exceedingly potent mediator of new blood vessel formation. It hydrolyzes cellular tRNAs resulting in decreased protein synthesis and is similar to pancreatic ribonuclease. In addition, the mature peptide has antimicrobial activity against some bacteria and fungi, including *S. pneumoniae* and *C. albicans*. Alternative splicing results in two transcript variants encoding the same protein. This gene and the gene that encodes ribonuclease, RNase A family, 4 share promoters and 5' exons. Each gene splices to a unique downstream exon that contains its complete coding region. [provided by RefSeq, Aug 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using ANG I Monoclonal Antibody