



CLLD7 Monoclonal Antibody

Catalog No	YP-mAb-16149
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	RCBTB1
Immunogen	The antiserum was produced against synthesized peptide derived from human RCBTB1. AA range:251-300
Specificity	CLLD7 Monoclonal Antibody detects endogenous levels of CLLD7 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CLLD7;CLLL7;E4.5;GLP;RCBTB1;RP11 185C18.1;RCC1 and BTB domain-containing protein 1;Chronic lymphocytic leukemia deletion region gene 7 protein;CLL deletion region gene 7 protein;Regulator of chromosome condensation and BTB domain-containing protein 1
Observed Band	58kD
Calculated Molecular Weight	58kD
Cell Pathway	Nucleus .
Tissue Specificity	Ubiquitously expressed (PubMed:11306461, PubMed:27486781). In the retina, present in the nerve fiber layer and to a lesser extent in the inner and outer plexiform layers (at protein level) (PubMed:27486781).
Function	function:May be involved in cell cycle regulation by chromatin remodeling., similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 2 BTB (POZ) domains.,similarity:Contains 6 RCC1 repeats.,tissue specificity:Ubiquitously expressed.,



Background

This gene encodes a protein with an N-terminal RCC1 domain and a C-terminal BTB (broad complex, tramtrack and bric-a-brac) domain. In rat, over-expression of this gene in vascular smooth muscle cells induced cellular hypertrophy. In rat, the C-terminus of RCBTB1 interacts with the angiotensin II receptor-1A. In humans, this gene maps to a region of chromosome 13q that is frequently deleted in B-cell chronic lymphocytic leukemia and other lymphoid malignancies. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using CLLD7 Monoclonal Antibody