



Gα olf Monoclonal Antibody

Catalog No	YP-mAb-16173
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	GNAL
Immunogen	The antiserum was produced against synthesized peptide derived from human GNAL. AA range:41-90
Specificity	Gα olf Monoclonal Antibody detects endogenous levels of Gα olf protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Adenylate cyclase stimulating G alpha protein;olfactory type;Adenylate cyclase-stimulating G alpha protein;DYT25;G alpha olf;Gnal;GNAL_HUMAN;Guanine nucleotide binding protein (G protein);alpha activating activity polypeptide;alpha stimulating activity polypeptide;Guanine nucleotide binding protein G(olf) subunit alpha;Guanine nucleotide-binding protein G(olf) subunit alpha;Gα olf
Observed Band	40kD
Calculated Molecular Weight	40kD
Cell Pathway	plasma membrane,extracellular exosome,
Tissue Specificity	Detected in olfactory neuroepithelium, brain, testis, and to a lower extent in retina, lung alveoli, spleen. Trace amounts where seen in kidney, adrenal gland and liver. Found to be expressed in all the insulinomas examined.
Function	function:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems. G(olf) alpha mediates signal transduction within the olfactory neuroepithelium and the basal ganglia. May be involved in some aspect of visual transduction, and in mediating the effect of one or more hormones/neurotransmitters., similarity:Belongs to the G-alpha family. G(s) subfamily.,subunit:G proteins are



composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,tissue specificity:Detected in olfactory neuroepithelium, brain, testis, and to a lower extent in retina, lung alveoli, spleen. Trace amounts where seen in kidney, adrenal gland and liver. Found to be expressed in all the insulinomas examined.,

Background

G protein subunit alpha L(GNAL) Homo sapiens This gene encodes a stimulatory G protein alpha subunit which mediates odorant signaling in the olfactory epithelium. This protein couples dopamine type 1 receptors and adenosine A2A receptors and is widely expressed in the central nervous system. Mutations in this gene have been associated with dystonia 25 and this gene is located in a susceptibility region for bipolar disorder and schizophrenia. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Gα olf Monoclonal Antibody