



Lfc Monoclonal Antibody

Catalog No	YP-mAb-16181
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ARHGEF2
Immunogen	The antiserum was produced against synthesized peptide derived from in-ternal human ARHGEF2. AA range:383-432
Specificity	Lfc Monoclonal Antibody detects endogenous levels of Lfc protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AA48978;ARHG2;ARHG2_HUMAN;ARHGEF 2;ARHGEF-2;ARHGEF2;GEF; GEF H1;GEF-H1;GEFH1;Guanine nucleotide exchange factor H1;KIAA651; Lbcl1;Lfc;LFP4;MGC9568;Microtubule-regulated Rho-GEF;mKIAA651;P4; Proliferating cell nucleolar antigen p4;Protein GEF-H1;Rho guanine nucleotide exchange factor 2;rho/rac guanine nucleotide exchange factor (GEF) 2;rho/rac guanine nucleotide exchange factor 2;rho/rac guanine nucleotide exchange factor;KIAA0651;LFP40;Proliferating cell nucleolar antigen p40
Observed Band	101kD
Calculated Molecular Weight	101kD
Cell Pathway	Cytoplasm, cytoskeleton . Cytoplasm . Cell junction, tight junction . Golgi apparatus . Cytoplasm, cytoskeleton, spindle . Cell projection, ruffle membrane . Cytoplasmic vesicle . Localizes to the tips of cortical microtubules of the mitotic spindle during cell division, and is further released upon microtubule depolymerization (PubMed:15827085). Recruited into membrane ruffles induced by S.flexneri at tight junctions of polarized epithelial cells (PubMed:19043560). Colocalized with NOD2 and RIPK2 in vesicles and with the cytoskeleton (PubMed:21887730). .
Tissue Specificity	Brain,Cervix carcinoma,Epithelium,Platelet,



Function

domain:The DH (DBL-homology) domain interacts with and promotes loading of GTP on RhoA.,domain:The PH (pleckstrin-homology) domain is involved in microtubule binding and targeting to tight junctions.,function:Activates Rho-GTPases by promoting the exchange of GDP for GTP. May be involved in epithelial barrier permeability, cell motility and polarization, dendritic spine morphology, antigen presentation, leukemic cell differentiation, cell cycle regulation, and cancer. Binds Rac-GTPases, but does not seem to promote nucleotide exchange activity toward Rac-GTPases, which was uniquely reported in PubMed:9857026. May stimulate instead the cortical activity of Rac. Inactive toward CDC42, TC10, or Ras-GTPases.,online information:ARHGEF2 entry, PTM:Phosphorylation of Ser-886 by PAK1 induces binding to protein 14-3-3 zeta, promoting its relocation to microtubules and the inhibition of its activit

Background

Rho GTPases play a fundamental role in numerous cellular processes that are initiated by extracellular stimuli that work through G protein coupled receptors. The encoded protein may form complex with G proteins and stimulate rho-dependent signals. Alternatively spliced transcript variants encoding different isoforms have been identified.[provided by RefSeq, Jun 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Lfc Monoclonal Antibody