



AChRα 5 Monoclonal Antibody

Catalog No	YP-mAb-16371
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CHRNA5
Immunogen	The antiserum was produced against synthesized peptide derived from human AChRα5. AA range:166-215
Specificity	AChRα5 Monoclonal Antibody detects endogenous levels of AChRα5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Acetylcholine receptor nicotinic alpha 5 (neuronal);ACHA5_HUMAN;AChR;Cholinergic receptor neuronal nicotinic alpha polypeptide 5;Cholinergic receptor nicotinic alpha 5;Cholinergic receptor nicotinic alpha polypeptide 5;CHRNA 5;CHRNA5;LNCR2;NACHRA 5;NACHRA5;Neuronal acetylcholine receptor subunit alpha 5;Neuronal acetylcholine receptor subunit alpha-5;Neuronal nicotinic acetylcholine receptor alpha 5 subunit;Neuronal nicotinic acetylcholine receptor;alpha5 subunit;OTTHUMP18519;AChRα5
Observed Band	53kD
Calculated Molecular Weight	53kD
Cell Pathway	Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Eye,
Function	disease:Genetic variations in CHRNA5 may be associated with susceptibility to lung cancer type 2 (LNCR2) [MIM:612052].,function:After binding acetylcholine, the AChR responds by an extensive change in conformation that affects all subunits and leads to opening of an ion-conducting channel across the plasma membrane.,similarity:Belongs to the ligand-gated ionic channel (TC 1.A.9)



family.,subunit:Neuronal AChR seems to be composed of two different type of subunits: alpha and non-alpha (beta).,

Background

The protein encoded by this gene is a nicotinic acetylcholine receptor subunit and a member of a superfamily of ligand-gated ion channels that mediate fast signal transmission at synapses. These receptors are thought to be heteropentamers composed of separate but similar subunits. Defects in this gene have been linked to susceptibility to lung cancer type 2 (LNCR2).[provided by RefSeq, Jun 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using AChRα 5 Monoclonal Antibody