



GABAA R α 6 Monoclonal Antibody

Catalog No	YP-mAb-16416
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	GABRA6
Immunogen	The antiserum was produced against synthesized peptide derived from human GABRA6. AA range:61-110
Specificity	GABAA R α 6 Monoclonal Antibody detects endogenous levels of GABAA R α 6 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ECA5;GABA alpha receptor beta-2 subunit;GABA(A) receptor subunit beta-3; GABAA receptor beta 3 subunit;GABAA receptor subunit beta 3;GABR B3; Gabrb3;Gamma aminobutyric acid (GABA) A receptor beta 3;Gamma aminobutyric acid receptor subunit beta 3;Gamma-aminobutyric acid receptor subunit beta-3;GBRB3_HUMAN;MGC951;GABAA R α 6;GABRA6;Gamma-aminobutyric acid receptor subunit alpha-6;GABA(A) receptor subunit alpha-6
Observed Band	51kD
Calculated Molecular Weight	51kD
Cell Pathway	Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Cerebellum,
Function	function:GABA, the major inhibitory neurotransmitter in the vertebrate brain, mediates neuronal inhibition by binding to the GABA/benzodiazepine receptor and opening an integral chloride channel.,online information:Forbidden fruit - Issue 56 of March 2005,similarity:Belongs to the ligand-gated ionic channel (TC 1.A.9) family.,subunit:Binds UBQLN1 (By similarity). Generally pentameric.



There are five types of GABA(A) receptor chains: alpha, beta, gamma, delta, and rho.,

Background

GABA is the major inhibitory neurotransmitter in the mammalian brain where it acts at GABA-A receptors, which are ligand-gated chloride channels. Chloride conductance of these channels can be modulated by agents such as benzodiazepines that bind to the GABA-A receptor. At least 16 distinct subunits of GABA-A receptors have been identified. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using GABAA R α 6 Monoclonal Antibody