



Cdc37L1 Monoclonal Antibody


Catalog No	YP-mAb-16684
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CDC37L1
Immunogen	Synthesized peptide derived from Cdc37L1 . at AA range: 80-160
Specificity	Cdc37L1 Monoclonal Antibody detects endogenous levels of Cdc37L1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CDC37B;CDC37L1;HARC;Hsp90 co-chaperone Cdc37-like 1;Hsp90-associating relative of Cdc37
Observed Band	38kD
Calculated Molecular Weight	38kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Expressed in brain, heart, kidney, liver, placenta and skeletal muscle.
Function	function:Co-chaperone that binds to numerous proteins and promotes their interaction with Hsp70 and Hsp90.,sequence caution:Translated as Cys., similarity:Belongs to the CDC37 family.,subunit:Self-associates. Forms complexes with Hsp70 and Hsp90. Interacts with CDC37, FKBP4, PPID and STIP1.,tissue specificity:Expressed in brain, heart, kidney, liver, placenta and skeletal muscle.,
Background	CDC37L1 is a cytoplasmic phosphoprotein that exists in complex with HSP90 (HSPCA; MIM 140571) as well as several other proteins involved in HSP90-mediated protein folding (Scholz et al., 2001 [PubMed 11413142]).[supplied by OMIM, Mar 2008],

**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using Cdc37L1 Monoclonal Antibody