



CUL-2 Monoclonal Antibody

Catalog No	YP-mAb-16718
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CUL2
Immunogen	The antiserum was produced against synthesized peptide derived from human Cullin 2. AA range:696-745
Specificity	CUL-2 Monoclonal Antibody detects endogenous levels of CUL-2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CUL 2;CUL2;cullin 2;CUL-2;Cullin-2
Observed Band	87kD
Calculated Molecular Weight	87kD
Cell Pathway	nucleoplasm,nucleolus,cytosol,VCB complex,cullin-RING ubiquitin ligase complex,Cul2-RING ubiquitin ligase complex,
Tissue Specificity	Brain,Kidney,Skin,
Function	function:Core component of multiple cullin-RING-based ECS (ElonginB/C-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of target proteins. May serve as a rigid scaffold in the complex and may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (By similarity). The functional specificity of the ECS complex depends on the substrate recognition component. ECS(VHL) mediates the ubiquitination of hypoxia-inducible factor (HIF).,pathway:Protein modification; protein ubiquitination.,PTM:CBC(VHL) complex formation seems to promote



neddylated. Deneddylated via its interaction with the COP9 signalosome (CSN

Background

function:Core component of multiple cullin-RING-based ECS (ElonginB/C-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of target proteins. May serve as a rigid scaffold in the complex and may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (By similarity). The functional specificity of the ECS complex depends on the substrate recognition component. ECS(VHL) mediates the ubiquitination of hypoxia-inducible factor (HIF).,pathway:Protein modification; protein ubiquitination.,PTM:CBC(VHL) complex formation seems to promote neddylation. Deneddylated via its interaction with the COP9 signalosome (CSN) complex.,similarity:Belongs to the cullin family.,subunit:Component of multiple ECS (Elongin BC-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes formed of CUL2, Elongin BC (TCEB1 and TCEB2), RBX1 and a variable substrate-specific adapter. Component of the ECS(VHL) or CBC(VHL) complex containing VHL. Component of the ECS(MED8) complex with the probable substrate recognition component MED8 (By similarity). Component of the ECS(PPIL5) complex with the probable substrate recognition component PPIL5. Component of a probable ECS E3 ubiquitin-protein ligase complex containing CUL2, RBX1, TCEB1, TCEB2 and FEM1B. Part of an E3 ubiquitin-protein ligase complex including ZYG11B, CUL2 and Elongin BC. Part of an E3 ubiquitin-protein ligase complex including ZYG11BL, CUL2 and Elongin BC.Interacts with RBX1, RNF7, FEM1B and TIP120A/CAND1. Interacts with COPS2, and MED8 (By similarity). Interacts with human respiratory syncytial virus (HRSV) protein NS1.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using CUL-2 Monoclonal Antibody