



RFC1 Monoclonal Antibody

Catalog No	YP-mAb-16793
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	RFC1
Immunogen	Synthesized peptide derived from the Internal region of human RFC1.
Specificity	RFC1 Monoclonal Antibody detects endogenous levels of RFC1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DNA binding protein PO GA;Activator 1 subunit 1;Activator 1 large subunit; Activator 1 140 kDa subunit;A1 140 kDa subunit;RFC1;Replication factor C subunit 1;RFC140;DNA-binding protein PO-GA;Replication factor C 140 kDa subunit;RF-C 140 kDa subunit;Replication factor C large subunit
Observed Band	128kD
Calculated Molecular Weight	128kD
Cell Pathway	Nucleus.
Tissue Specificity	Wide tissue distribution. Undetectable in placental tissue.
Function	function:Interacts with C-terminus of PCNA. 5' phosphate residue is required for binding of the N-terminal DNA-binding domain to duplex DNA, suggesting a role in recognition of non-primer template DNA structures during replication and/or repair.,function:The elongation of primed DNA templates by DNA polymerase delta and epsilon requires the action of the accessory proteins PCNA and activator 1. This subunit binds to the primer-template junction. Binds the PO-B transcription element as well as other GA rich DNA sequences. Could play a role in DNA transcription regulation as well as DNA replication and/or repair. Can bind single- or double-stranded DNA.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the activator 1 large



subunit family.,similarity:Contains 1 BRCT domain.,subunit:Heterotetramer of subunits RFC2, RFC3, RFC4 and RFC5 that can form a complex

Background

This gene encodes the large subunit of replication factor C, a five subunit DNA polymerase accessory protein, which is a DNA-dependent ATPase required for eukaryotic DNA replication and repair. The large subunit acts as an activator of DNA polymerases, binds to the 3' end of primers, and promotes coordinated synthesis of both strands. It may also have a role in telomere stability. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq, Mar 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using RFC1 Monoclonal Antibody