



Cleaved-Integrin α V HC (K889) Monoclonal Antibody

Catalog No	YP-mAb-16808
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ITGAV
Immunogen	The antiserum was produced against synthesized peptide derived from human ITGAV. AA range:840-889
Specificity	Cleaved-Integrin α V HC (K889) Monoclonal Antibody detects endogenous levels of fragment of activated Integrin α V HC protein resulting from cleavage adjacent to K889.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	antigen identified by monoclonal antibody L23;CD 51;CD51;DKFZp686A8142; integrin alpha V beta 3;integrin alpha-V;Integrin alpha-V light chain;integrin alphaVbeta3;integrin;alpha V (vitronectin receptor;alpha polypeptide;antigen CD51);ITAV_HUMAN;ITGAV;MSK 8;Msk8;Vitronectin receptor subunit alpha; VNRA;VTNR;Integrin α V;CD antigen CD51
Observed Band	95kD
Calculated Molecular Weight	95kD
Cell Pathway	Cell membrane; Single-pass type I membrane protein. Cell junction, focal adhesion .
Tissue Specificity	Aortic endothelium,Liver,Pooled,Testis,
Function	function:The alpha-V integrins are receptors for vitronectin, cytotactin, fibronectin, fibrinogen, laminin, matrix metalloproteinase-2, osteopontin, osteomodulin, prothrombin, thrombospondin and vWF. They recognize the sequence R-G-D in a wide array of ligands. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in



Kaposi's sarcoma lesions.,similarity:Belongs to the integrin alpha chain family.,similarity:Contains 7 FG-GAP repeats.,subunit:Heterodimer of an alpha and a beta subunit. The alpha subunit is composed of an heavy and a light chain linked by a disulfide bond. Alpha-V associates with either beta-1, beta-3, beta-5, beta-6 or beta-8 subunit. Interacts with HIV-1 Tat. Alpha-V/beta-6 binds to foot-and-mouth disease virus (FMDV) VP1 protein and acts as a receptor for this virus (By similarity). Alpha-V/beta-6 binds to coxsack

Background

integrin subunit alpha V(ITGAV) Homo sapiens The product of this gene belongs to the integrin alpha chain family. Integrins are heterodimeric integral membrane proteins composed of an alpha subunit and a beta subunit that function in cell surface adhesion and signaling. The encoded preproprotein is proteolytically processed to generate light and heavy chains that comprise the alpha V subunit. This subunit associates with beta 1, beta 3, beta 5, beta 6 and beta 8 subunits. The heterodimer consisting of alpha V and beta 3 subunits is also known as the vitronectin receptor. This integrin may regulate angiogenesis and cancer progression. Alternative splicing results in multiple transcript variants. Note that the integrin alpha 5 and integrin alpha V subunits are encoded by distinct genes. [provided by RefSeq, Oct 2015],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Cleaved-Integrin α V HC (K889) Monoclonal Antibody