



K-cadherin Monoclonal Antibody

Catalog No	YP-mAb-17078
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CDH6
Immunogen	Synthesized peptide derived from K-cadherin . at AA range: 180-260
Specificity	K-cadherin Monoclonal Antibody detects endogenous levels of K-cadherin protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CAD6;KCAD;K-Cadherin (CDH6);K-cadherin;CDH6;Cadherin-6;Kidney cadherin
Observed Band	88kD
Calculated Molecular Weight	88kD
Cell Pathway	Cell membrane; Single-pass type I membrane protein.
Tissue Specificity	Highly expressed in brain, cerebellum, and kidney. Lung, pancreas, and gastric mucosa show a weak expression. Also expressed in certain liver and kidney carcinomas.
Function	function:Cadherins are calcium dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types., similarity:Contains 5 cadherin domains.,tissue specificity:Highly expressed in brain, cerebellum, and kidney. Lung, pancreas, and gastric mucosa show a weak expression. Also expressed in certain liver and kidney carcinomas.,
Background	This gene encodes a member of the cadherin superfamily. Cadherins are membrane glycoproteins that mediate homophilic cell-cell adhesion and play critical roles in cell differentiation and morphogenesis. The encoded protein is a



type II cadherin and may play a role in kidney development as well as endometrium and placenta formation. Decreased expression of this gene may be associated with tumor growth and metastasis. [provided by RefSeq, May 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using K-cadherin Monoclonal Antibody