



Frizzled-8 Monoclonal Antibody

Catalog No	YP-mAb-13262
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	FZD8
Immunogen	The antiserum was produced against synthesized peptide derived from human FZD8. AA range:486-535
Specificity	Frizzled-8 Monoclonal Antibody detects endogenous levels of Frizzled-8 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Frizzled 8;FZ 8;FZD8;hFZ8;Frizzled-8;Fz-8
Observed Band	70kD
Calculated Molecular Weight	70kD
Cell Pathway	Membrane; Multi-pass membrane protein. Golgi apparatus . Cell membrane ; Multi-pass membrane protein . Colocalizes with GOPC at the Golgi apparatus. .
Tissue Specificity	Most abundant in fetal kidney, followed by brain and lung. In adult tissues, expressed in kidney, heart, pancreas and skeletal muscle.
Function	domain:Lys-Thr-X-X-X-Trp motif is involved in the activation of the Wnt/beta-catenin signaling pathway.,domain:The FZ domain is involved in binding with Wnt ligands.,domain:The PDZ-binding motif mediates interaction with GOPC.,function:Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes has been seen for some family members, but it is not yet clear if it represents a distinct pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for Wnt-mediated inactivation



of GSK-3 kinase. Both pathways seem to involve interactions with G-proteins. May be involved in transduction and intercellul

Background

frizzled class receptor 8(FZD8) Homo sapiens This intronless gene is a member of the frizzled gene family. Members of this family encode seven-transmembrane domain proteins that are receptors for the Wingless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. This gene is highly expressed in two human cancer cell lines, indicating that it may play a role in several types of cancer. The crystal structure of the extracellular cysteine-rich domain of a similar mouse protein has been determined. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using Frizzled-8 Monoclonal Antibody

