



PCNA mouse mAb

Catalog No	YP-Ab-16574
Isotype	IgG
Reactivity	Human;Mouse;Monkey;Rat;Hamster
Applications	WB;IP;ICC;IHC
Gene Name	pcna
Immunogen	Purified recombinant human PCNA protein fragments expressed in E.coli.
Specificity	This antibody detects endogenous levels of PCNA and does not cross-react with related proteins.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	wb dilution 1:1000 icc dilution 1:100
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Cyclin;proliferating cell nuclear antigen;PCNA;DNA polymerase delta auxiliary protein;HGCN8729;MGC8367;Mutagen-sensitive 209 protein; OTTHUMP00000030189;OTTHUMP00000030190;Pcna/cyclin;PCNA_HUMAN; PCNAR;Polymerase delta accessory protein;Proliferating Cell Nuclear Antigen.
Observed Band	30-33kd
Calculated Molecular Weight	30-33kd
Cell Pathway	Nucleus . Colocalizes with CREBBP, EP300 and POLD1 to sites of DNA damage (PubMed:24939902). Forms nuclear foci representing sites of ongoing DNA replication and vary in morphology and number during S phase (PubMed:15543136). Co-localizes with SMARCA5/SNF2H and BAZ1B/WSTF at replication foci during S phase (PubMed:15543136). Together with APEX2, is redistributed in discrete nuclear foci in presence of oxidative DNA damaging agents. .
Tissue Specificity	Bone marrow,Fetal brain cortex,Lung,Placenta,
Function	disease:Antibodies are present in sera from patients with systemic lupus erythematosus.,function:This protein is an auxiliary protein of DNA polymerase delta and is involved in the control of eukaryotic DNA replication by increasing




the polymerase's processibility during elongation of the leading strand.,online information:PCNA entry,PTM:Upon methyl methanesulfonate-induced DNA damage, mono-ubiquitinated by the UBE2B-RAD18 complex on Lys-164. This induces non-canonical poly-ubiquitination on Lys-164 through 'Lys-63' linkage of ubiquitin moieties by the E2 complex UBE2N-UBE2V2 and the E3 ligases RNF8 and SHPRH, which are required for DNA repair.,similarity:Belongs to the PCNA family.,subunit:Homotrimer. Interacts with KCTD10. Interacts with PPP1R15A (By similarity). Forms a complex with activator 1 heteropentamer in the presence of ATP. Interacts with POLH, POLK, DNMT1, ERCC5/XPG, FEN1, C


Background	The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images




Western blot detection of PCNA in HeLa, 3T3, COS7, C6 and CHO-K1 cell lysates using PCNA mouse mAb (1:1000 diluted). Predicted band size: 36KDa. Observed band size: 36KDa.



Immunocytochemistry staining of HeLa cells using PCNA mouse mAb (dilution 1:100). Fixed in 100% methanol for 2hr at -20°C.



Immunoprecipitation analysis of Hela cell lysates using PCNA mouse mAb.



Immunohistochemical analysis of paraffin-embedded human colorectal carcinoma with PCNA mouse mAb (2E1-G10-H10, 1:400 diluted), showing nuclear localization. A high pressure mediated antigen retrieval step was performed in citrate buffer (pH 6.0).