



PI 3 kinase p85/p55 (phospho Tyr467/199) Polyclonal Antibody

货号	YP-Ab-17854
同位型	IgG
应用	WB;IHC;IF;ELISA
种属	Human;Mouse;Rat;Monkey;Pig
靶点	PI3 kinase p85/p55
简介	>>EGFR tyrosine kinase inhibitor resistance;>>Endocrine resistance;>>Platinum drug resistance;>>ErbB signaling pathway;>>Ras signaling pathway;>>Rap1 signaling pathway;>>cAMP signaling pathway;>>Chemokine signaling pathway;>>HIF-1 signaling pathway;>>FoxO signaling pathway;>>Phosphatidylinositol signaling system;>>Sphingolipid signaling pathway;>>Phospholipase D signaling pathway;>>Autophagy - animal;>>mTOR signaling pathway;>>PI3K-Akt signaling pathway;>>AMPK signaling pathway;>>Apoptosis;>>Longevity regulating pathway;>>Longevity regulating pathway - multiple species;>>Cellular senescence;>>Axon guidance;>>VEGF signaling pathway;>>Osteoclast differentiation;>>Focal adhesion;>>Signaling pathways regulating pluripotency of stem cells;>>Platelet activation;>>Neutrophil extracellular trap formation;>>Toll-like receptor signaling pathway;>>C-type lectin receptor signaling pathway;>>JAK-STAT signaling pathway;>>Natural killer cell mediated cytotoxicity;>>T cell receptor signaling pathway;>
基因名称	PIK3R1/PIK3R3
免疫原	The antiserum was produced against synthesized peptide derived from human PI3-kinase p85-alpha/gamma around the phosphorylation site of Tyr467/199. AA range:436-485
特异性	Phospho-PI 3-kinase p85/p55 (Y467/199) Polyclonal Antibody detects endogenous levels of PI 3-kinase p85/p55 protein only when phosphorylated at Y467/199.
组成	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
来源	Polyclonal, Rabbit,IgG
稀释	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
纯化工艺	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
其他名称	GRB 1;GRB1;p85 alpha;p85;P85A_HUMAN;Phosphatidylinositol 3 kinase associated p 85 alpha;Phosphatidylinositol 3 kinase regulatory 1; Phosphatidylinositol 3 kinase regulatory subunit alpha;Phosphatidylinositol 3 kinase regulatory subunit polypeptide 1 (p85 alpha);Phosphatidylinositol 3-kinase 85 kDa regulatory subunit alpha;Phosphatidylinositol 3-kinase regulatory subunit alpha;Phosphoinositide 3 kinase regulatory subunit 1 (alpha); Phosphoinositide 3 kinase regulatory subunit 1 (p85 alpha);Phosphoinositide 3 kinase regulatory subunit 1;Phosphoinositide 3 kinase regulatory subunit







polypeptide 1 (p85 alpha);PI3 kinase p85 subunit alpha;PI3-kinase regulatory subunit alpha;PI3-kinase subunit p85-alpha;PI3K;PI3K regulatory subunit alpha;Pik3r1;PtdIns 3 kinase p85 alpha;PtdIns-3-kinase regulatory subunit alpha;PtdIns-3-kinase regulatory subunit p85-alpha;DKFZp686P5226;FLJ41892;OTTHUMP9783;OTTHUMP9786;p55;p55 gamma;P55G_HUMAN;p55PIK;Phosphatidylinositol 3 kinase regulatory subunit gamma;Phosphatidylinositol 3 kinase regulatory subunit polypeptide 3;Phosphatidylinositol 3 kinase;regulatory subunit;polypeptide 3 (p55;gamma);Phosphatidylinositol 3-kinase 55 kDa regulatory subunit gamma;Phosphatidylinositol 3-kinase regulatory subunit gamma;Phosphoinositide 3 kinase regulatory subunit 3 (gamma);Phosphoinositide 3 kinase regulatory subunit 3;Phosphoinositide 3 kinase regulatory subunit polypeptide 3;Phosphoinositide 3 kinase;regulatory subunit 3 (p55;PI3 kinase p85 subunit gamma;PI3-kinase regulatory subunit gamma;PI3-kinase subunit p55-gamma;PI3K regulatory subunit gamma;Pik3r3;PtdIns 3 kinase p85 gamma;PtdIns-3-kinase regulatory subunit gamma;PtdIns-3-kinase regulatory subunit p55-gamma;PI3 kinase p85/p55;PIK3R1/PIK3R3;Phosphatidylinositol 3-kinase regulatory subunit alpha/gamma

实测条带	55+85kD
分子量	54kDa,84kDa; 84kD,54kD(Calculated).
背景	Phosphatidylinositol 3-kinase phosphorylates the inositol ring of phosphatidylinositol at the 3-prime position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50 kD. This gene encodes the 85 kD regulatory subunit. Phosphatidylinositol 3-kinase plays an important role in the metabolic actions of insulin, and a mutation in this gene has been associated with insulin resistance. Alternative splicing of this gene results in four transcript variants encoding different isoforms. [provided by RefSeq, Jun 2011],
功能	disease:Defects in PIK3R1 are a cause of severe insulin resistance.,domain:The SH3 domain mediates the binding to CBLB, and to HIV-1 Nef.,function:Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues.,PTM:Polyubiquitinated in T-cells by CBLB; which does not promote proteasomal degradation but impairs association with CD28 and CD3Z upon T-cell activation.,similarity:Belongs to the PI3K p85 subunit family.,similarity:Contains 1 Rho-GAP domain.,similarity:Contains 1 SH3 domain.,similarity:Contains 2 SH2 domains.,subunit:Heterodimer of a p110 (catalytic) and a p85 (regulatory) subunits. Interacts with phosphorylated TOM1L1. Interacts with phosphorylat
细胞定位	nucleus,cytoplasm,cis-Golgi network,cytosol,plasma membrane,cell-cell junction, phosphatidylinositol 3-kinase complex,phosphatidylinositol 3-kinase complex, class IA,membrane,perinuclear endoplasmic reticulum membrane,
组织表达	Isoform 2 is expressed in skeletal muscle and brain, and at lower levels in kidney and cardiac muscle. Isoform 2 and isoform 4 are present in skeletal muscle (at protein level).
浓度	1 mg/ml
储存	-15°C to -25°C/1 year(Do not lower than -25°C)
有关注意事项	Avoid repeated freezing and thawing!
使用建议	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images






A

B

C



A

B

C