DSS1 Rabbit pAb

Catalog No	YP-Ab-19571
Isotype	IgG
Reactivity	Human,Mouse
Applications	WB
Gene Name	SHFM1 DSS1 SHFDG1
Immunogen	Synthesized peptide derived from human DSS1. AA range:1-70
Specificity	This antibody detects endogenous levels of DSS1 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-15°C to -25°C/1 year(Do not lower than -25°C)
Synonyms	SHFM1 DSS1 SHFDG1;26S proteasome complex subunit DSS1 (Deleted in split hand/split foot protein 1) (Split hand/foot deleted protein 1) (Split hand/foot malformation type 1 protein)
Observed Band	
Calculated Molecular Weight	8kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in limb bud , craniofacial primordia and skin.
Function	Component of the 26S proteasome , a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins , which could impair cellular functions , and by removing proteins whose functions are no longer required. Therefore , the proteasome participates in numerous cellular processes , including cell cycle progression , apoptosis , or DNA damage repair . Component of the TREX-2 complex (transcription and export complex 2) , composed of at least ENY2 , GANP , PCID2 , SEM1 , and either centrin CETN2 or CETN3 . The TREX-2 complex functions in docking export-competent ribonucleoprotein particles (mRNPs) to the nuclear entrance of the nuclear pore complex (nuclear basket) . TREX-2 participates in mRNA export and accurate chromatin positioning in the nucleus by tethering genes to the nuclear periphery. Binds and stabilizes BRCA2 and is thus involved

	in the control of R-loop-associated DNA damage and thus transcription-associated genomic instability. R-loop accumulation increases in SEM1-depleted cells.
Background	
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images