
# TOR3A mouse mAb

| | |
|---|---|
| **Catalog No** | YP-mAb-11772 |
| **Isotype** | IgG |
| **Reactivity** | Human;Mouse;Rat |
| **Applications** | WB |
| **Gene Name** | TOR3A ADIR |
| **Immunogen** | Synthesized peptide derived from human TOR3A AA range: 151-201 |
| **Specificity** | This antibody detects endogenous levels of TOR3A at Human/Mouse/Rat |
| **Formulation** | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| **Source** | Monoclonal, Mouse,IgG |
| **Purification** | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| **Dilution** | WB 1:500-1:2000 |
| **Concentration** | 1 mg/ml |
| **Purity** | ≥90% |
| **Storage Stability** | -20°C/1 year |
| **Synonyms** | torsin family 3;member A;Torsin family 3 member A;Torsin 3A;ATP-dependent interferon-responsive protein;ADIR2;TOR3A;TOR3A ADIR |
| **Observed Band** | |
| **Calculated Molecular Weight** | 44kD |
| **Cell Pathway** | Cytoplasm. Endoplasmic reticulum lumen. |
| **Tissue Specificity** | Ubiquitously expressed. Highest expression in stomach, salivary glands and lymph nodes. Isoform 2 is expressed in placenta. |
| **Function** | similarity:Belongs to the clpA/clpB family. Torsin subfamily.,tissue specificity:Ubiquitously expressed. Highest expression in stomach, salivary glands and lymph nodes. Isoform 2 is expressed in placenta., |
| **Background** | |
| **matters needing attention** | Avoid repeated freezing and thawing! |
| **Usage suggestions** | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

## Products Images




Western Blot analysis of various cells using TOR3A mouse mAb